

MODERNA COVID-19 VACCINATION SCHEDULE AND DOSING

FOR NON-IMMUNOCOMPROMISED POPULATIONS


Moderna

Ages 6 months–5 years

dose/injection volume
(Do NOT dilute before use)

Primary Dose:

Blue Cap (magenta label)



Moderna

Ages 6 years–11 years

dose/injection volume
(Do NOT dilute before use)

Primary Dose:

Blue Cap (purple label)



Moderna

Ages 12 years–17 years

dose/injection volume
(Do NOT dilute before use)

Primary Dose:

Red Cap (blue label)

PFIZER ONLY Bivalent Booster:

Gray Cap (gray label)



Moderna

Ages 18 years and older

dose/injection volume
(Do NOT dilute before use)


Primary Dose:

Red Cap (blue label)

Bivalent Booster:

MODERNA Dark Blue Cap (gray label) or

PFIZER Gray Cap (gray label)



* Complete the primary series with same product. If the vaccine product previously administered cannot be determined or is no longer available, any age-appropriate mRNA COVID-19 vaccine product may be administered at least 28 days after the first dose.

† Persons with a recent SARS-CoV-2 infection may consider delaying a primary series or booster dose by 3 months from symptom onset or positive test (if infection was asymptomatic).

‡ Some studies in adolescents and adults have shown the small risk of myocarditis associated with mRNA COVID-19 vaccines might be reduced and peak antibody responses and vaccine effectiveness may be increased with an interval longer than 4 weeks. An 8-week interval may be optimal for people who are not moderately or severely immunocompromised and ages 6 months–64 years, especially for males ages 12–39 years. Source: [CDC](https://www.cdc.gov/media/releases/2022/s0927-covid19-vaccine-interval.html).



See the [Moderna COVID-19 Vaccine Guidance](#)

for children who are transitioning from a younger to older age group during their vaccination window.

See the [CDC's Guidance](#) for latest updates and information on who is considered moderately or severely immunocompromised.