


OTHER COVID-19 VACCINATION SCHEDULE AND DOSING

Janssen (J&J) Ages 18 years and older

dose/injection volume
(dilute before use)

Bivalent Boosters:

Pfizer Gray Cap (gray label) or
Moderna Dark Blue Cap (gray label)



Novavax Ages 12 years and older


dose/injection volume

Bivalent Booster Options:

Pfizer Gray Cap (gray label)
Moderna Dark Blue Cap (gray label)

Monovalent Booster:

Novavax Light Gray Cap¹



¹ Per the [CDC](#): A single Novavax booster dose (instead of a bivalent mRNA booster dose) may be given to persons 18 years of age or older who have not received a previous booster dose in limited situations. These situations are 1. an mRNA vaccine is contraindicated, or not available or 2. the recipient is unwilling to receive an mRNA vaccine and would otherwise not receive a booster dose. Administer the booster dose at least 6 months after the last primary series dose.

FOR MODERATELY TO SEVERELY IMMUNOCOMPROMISED POPULATIONS

Janssen (J&J) Ages 18 years and older


dose/injection volume
(dilute before use)

Primary Dose:

Pfizer's Gray Cap
Moderna's Red Cap (blue label)

Bivalent Boosters:

Pfizer Gray Cap (gray label) or
Moderna Dark Blue Cap (gray label)



§ Age-appropriate mRNA COVID-19 vaccines are preferred over the Janssen COVID-19 Vaccine for all vaccine doses for all vaccine-eligible people 18 years of age and older. Janssen COVID-19 vaccine should only be used in limited situations. See [CDC](#).

¶ mRNA vaccine must be used.

Source: [COVID-19 Vaccine Interim COVID-19 Immunisation Schedule for 6 Months of Age and Older](#)

See the [CDC's Guidance](#) for latest updates and information on who is considered moderately or severely immunocompromised.

Funding for this project was made possible by the Office of Disease Control, through the Illinois Department of Public Health.